

เอกสารกำกับยาภาษาอังกฤษ

DIPHTHERIA AND TETANUS VACCINE ADSORBED FOR ADULTS AND ADOLESCENTS

DESCRIPTION

Diphtheria and tetanus vaccine adsorbed for adults and adolescents (Td) is prepared by combining purified diphtheria toxoid and purified tetanus toxoid. The antigens are adsorbed onto Aluminium Phosphate as adjuvant. Thiomersal is added as preservative. The vaccine has the appearance of a greyish – white suspension and does not contain any horse serum protein. Therefore, it does not induce sensitization to sera of equine origin. The vaccine meets the requirements of W.H.O. and B.P. when tested by the methods outlined in W.H.O., TRS. 980 (2014) and B.P.

POTENCY

Each single 0.5 ml human dose contains

Diphtheria Toxoid ≤ 5 Lf (≥ 2 IU)

Tetanus Toxoid ≥ 5 Lf (≥ 40 IU)

Adsorbed on Aluminium Phosphate, $Al^{+++} \leq 1.25$ mg

Preservative: 0.005% Thiomersal

INDICATIONS

For primary vaccination and revaccination of adults and adolescents, who are having contraindications of DTP.

Primary vaccination and revaccination of children older than 7 years. In order to prevent allergic reactions to the protein of Diphtheria toxoid, the quantity of the toxoid has been markedly reduced.

After a primary immunization course of either DTP or Td, adsorbed Td for adults may be used as a booster at intervals of approximately 10 years, but with a minimum of at least one year between doses. It can safely replace monovalent tetanus toxoid (TT) vaccine, including during pregnancy.

The vaccine can be safely and effectively given simultaneously with BCG, Measles, Polio Vaccine (IPV and OPV), Hepatitis B, Yellow fever Vaccine, *Haemophilus influenzae* type b, Varicella vaccine and vitamin A supplementation.

APPLICATION AND DOSAGE

Two injections of 0.5 ml at least four weeks apart followed by a third injection 6 to 12 months after the second dose. The vaccine should also be given as a booster immunization every 5 to 10 years.

METHOD OF INNOCULATION

The vaccine should be injected intramuscularly. The preferred site for injection is deltoid muscles. Care should be taken not to inject into the blood vessel or the skin. Only sterile syringes and needles should be used for each injection. The vaccine should be well shaken before use.

Once opened, multi-dose vials should be kept between +2°C and +8°C. Multi-dose vials of Diphtheria and tetanus vaccine adsorbed for adults and adolescents vaccine from which one or more doses of vaccine have been removed during an immunization session may be used in subsequent immunization sessions for upto a maximum of 28 days, provided that all of the following conditions are met (as described in the WHO policy statement: Handling of multi dose vaccine vials after opening, WHO/IVB/14.07):

- The vaccine is currently prequalified by WHO;
- The vaccines is approved for use for up to 28 days after opening the vial, as determined by WHO;
- The expiry date of the vaccine has not passed;
- The vaccine vial has been, and will continue to be, store at WHO- or manufacturer recommended temperatures; furthermore, the vaccine vial monitor, if one is attached, is visible on the vaccine label and is not past its discard point, and the vaccine has not been damaged by freezing.

The vaccine should be visually inspected for any foreign particulate matter and / or variation of physical aspect prior to administration. In event of either being observed, discard the vaccine.

REACTIONS

Reactions are generally mild and confined to the site of injection. Some inflammation may occur together with transient fever, malaise and irritability. Occasionally a nodule may develop at the site of injection but this is rare.

PRECAUTIONS

ADRENALINE INJECTION (1:1000) MUST BE IMMEDIATELY AVAILABLE SHOULD AN ACUTE ANAPHYLACTIC REACTION OCCUR DUE TO ANY COMPONENT OF THE VACCINE. For treatment of severe anaphylaxis the initial dose of adrenaline is 0.1-0.5 mg (0.1-0.5 ml of 1:1000 injection) given S/C or I/M. Single dose should not exceed 1 mg (1 ml). For infants and children the recommended dose of adrenaline is 0.01 mg/kg (0.01 ml/kg of 1:1000 injection). Single paediatric dose should not exceed 0.5 mg (0.5 ml). The mainstay in the treatment of severe anaphylaxis is the prompt use of adrenaline, which can be lifesaving. It should be used at the first suspicion of anaphylaxis.

As with the use of all vaccines the vaccine should remain under observation for not less than 30 minutes for possibility of occurrence of immediate or early allergic reactions.

Hydrocortisone and antihistaminics should also be available in addition to such as oxygen inhalation.

Special care should be taken to ensure that the injection does not enter a blood vessel.

IT IS EXTREMELY IMPORTANT WHEN THE PARENT, GUARDIAN, OR ADULT PATIENT RETURNS FOR THE NEXT DOSE IN THE SERIES, THE PARENT, GUARDIAN, OR ADULT PATIENT SHOULD BE QUESTIONED CONCERNING OCCURRENCE OF ANY SYMPTOMS AND / OR SIGNS OR AN ADVERSE REACTION AFTER THE PREVIOUS DOSE.

DRUG INTERACTIONS

If Td and TIG or Diphtheria Antitoxin are administered concurrently, separate syringes and separate sites should be used.

As with other intramuscular injections, use with caution in patients on anticoagulant therapy. Immunosuppressive therapies, including irradiation, antimetabolites, alkylating agents, cytotoxic drugs, and corticosteroids used in greater than physiologic doses), may reduce the immune response to vaccines.

CONTRAINDICATIONS AND WARNINGS

The vaccine should not be given to persons who showed a severe reaction to a previous dose of Diphtheria and Tetanus vaccine.

A history of systemic allergic or neurologic reactions following a previous dose of Td is an absolute contraindication for further use.

Immunization should be deferred during the course of an acute illness. Vaccination of persons with severe, febrile illness should generally be deferred until these persons have recovered. However, the presence of minor illnesses such as mild upper respiratory infections with or without fever should not preclude vaccination.

IMMUNE DEFICIENCY

Individuals infected with human immunodeficiency virus (HIV), both asymptomatic and symptomatic, should be immunized with Td vaccine according to standard schedules.

STORAGE OF THE VACCINE

The vaccine should be stored in a dry, dark place at a temperature between 2-8°C.

Transportation should also be at 2-8°C. DO NOT FREEZE.

SHELF LIFE

36 months from date of manufacture.

PRESENTATION

1 dose ampoule of 0.5 ml

10 dose vial of 5 ml

20 dose vial of 10 ml

Manufactured by:

SERUM INSTITUTE OF INDIA PVT. LTD.

212/2, Hadapsar, Pune 411028, INDIA

Protection from birth onwards

นำส่งโดย

บริษัท มาสุ จำกัด

ทท. 02-556-1710-19